

Winter BB Cream OW I-17.932.01

Phase	Material Name	US INCI	Supplier	% Material
A	Montanov 68 MB	Cetearyl Alcohol Cetearyl Glucoside	SEPPIC	3.00
А	Montanov 202	Arachidyl Alcohol Behenyl Alcohol Arachidyl Glucoside	SEPPIC	3.00
A	Eutanol G	Octyldodecanol	BASF	2.00
A	Luvitol Lite EM	Hydrogenated Polyisobutene	BASF	5.00
A	Cutina FS 45	Stearic Acid Palmitic Acid	BASF	1.00
В	Deionized Water	Water	N.A.	66.63
В	Keltrol CG-RD	Xanthan Gum	CP Kelco	0.20
В	Prestige Soft Silver	Titanium Dioxide Mica Tin Oxide	Sudarshan	3.80
В	Colorona Bright Gold	Mica Titanium Dioxide Iron Oxides	Merck	3.00
В	Colorona Sienna Fine	Iron Oxides Mica	Merck	0.40
В	EDETA B Powder	Tetrasodium EDTA	BASF	0.10
С	Vitamin F forte	Linoleic Acid Linolenic Acid	CLR	1.00
С	Dermofeel Toco 70 non GMO	Tocopherol Helianthus Annuus (Sunflower) Seed Oil	Evonik	0.01
D	DayMoist CLR™	Water Hydrolyzed Corn Starch Beta Vulgaris (Beet) Root Extract	CLR	3.00
D	Phytosan™ K	Water Glycerin Glycine Soja (Soybean) Seed Extract	CLR	2.00
D	Microcare SB	Water Sodium Benzoate Potassium Sorbate	THOR	0.50
D	Agenaflo OS 9051	Aluminum Starch Octenylsuccinate	AGRANA	2.00
D	ProBarrier™ CLR	Water Caprylic/Capric Glycerides Copernicia Cerifera (Carnauba) Wax Decyl Glucoside Pentylene Glycol	CLR	3.00
E	Citric Acid (10%)	Water Citric Acid	N.A.	0.36
				100.00

Operating Instructions

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com



Winter BB Cream OW I-17.932.01

Mix A and predisperse phase B until completely uniform. Heat up A/B to 85°C separately. Add C to A and AC to B while stirring. Homogenize for 3 minutes and stir cold slowly. Add D in this order below 30°C and stir until homogeneous. Adjust pH value to 5-5.5 with F.

Directions for use:

Gently apply to face and neck for instant coverage and correction as well as for protection. Reapply as needed.

They are, to the best of our belief, correct, but are offered without obligation.